

# Impacts of environmental stress on carbon and water exchange of flatwoods pine forests in Florida

Timothy A. Martin

Greg Starr

Kenneth L. Clark


Thomas Powell

Henry L. Gholz

School of Forest Resources  
and Conservation,  
University of Florida




# Distribution of Florida Timberland



# Questions

- How do environmental stresses alter the carbon and water balance of flatwoods forests?
  - What sub-ecosystem scale processes are responsible for these responses?
  - At what time scales are ecosystem-, plant- and organ-level responses to stress expressed?



# Regenerating Clearcut



- Clearcut summer/fall 1997
- Double-bedded and planted winter 1998-1999
- Eddy covariance Feb 1998-present

# Mid-Rotation Plantation




- Established 1990
- Double-bedded,  
site prep herbicide,  
N&P @ 3 years
- Mean DBH = 9.8 cm
- Stand BA =  $16.1 \text{ m}^2 \text{ ha}^{-1}$
- Dominant height = 10 m
- $2032 \text{ trees ha}^{-1}$
- Peak all-sided LAI = 6.4

# Naturally Regenerated Stand



- *Pinus palustris / Pinus elliottii*
- Mean age ~ 60 years
- 333 stems  $\text{ha}^{-1}$
- Canopy height 22 m
- Mean tree DBH 25.3 cm
- Basal area  $18 \text{ m}^2 \text{ ha}^{-1}$
- All sided LAI
  - 2.6 summer
  - 2.0 winter

*Pinus elliottii*, Alachua County, Florida, 2000-2001




# Eddy Covariance



- Ecosystem-scale fluxes of
  - carbon
  - water
  - energy
- 3-D sonic anemometer
- Infrared gas analyzer

# AMERIFLUX System




- Long-term measurements of CO<sub>2</sub>, H<sub>2</sub>O and energy exchange in various ecosystems
- Collect information to help define the global C budget
- Enhance understanding of C fluxes, NEP and carbon sequestration


# Measurements

- Leaf-level
  - gas exchange
  - chl fluorescence
  - leaf water potential
  - nutrient concentration
- Tree-level
  - Granier sap flow probes
- Ecosystem-level
  - eddy covariance
  - PAR interception
  - soil CO<sub>2</sub> efflux
  - litterfall and other ecological meas.





Net Photosynthesis Rate





# Previous flatwoods research

- Intensive Management Practices Assessment Center (IMPAC)
  - early slash and loblolly pine growth not affected by irrigation




# Previous flatwoods research

- Teskey *et al.* 1994:
  - Slash pine net photosynthesis not impacted by water table drop < 1.5 m



# Previous southern pine research


- Albaugh *et al.* 1998:
  - Loblolly pine growing on a North Carolina sandhill is primarily nutrient limited





# Previous Research




- Cropper's Slash Pine Model successfully modeled carbon relations without water limitation functions







exchange ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )



January - March


$$\frac{0.0254 * \text{PPFD} * 26.56}{0.0254 * \text{PPFD} + 26.56}$$


April


$$\frac{0.0274 * \text{PPFD} * 24.72}{0.0274 * \text{PPFD} + 24.72}$$


May

$$\frac{0.0227 * \text{PPFD} * 10.18}{0.0227 * \text{PPFD} + 10.18}$$












# Hydraulic Conductance





Leaf Water Potential  
(MPa)

Tree Transpiration Rate  
( $\text{mol m}^{-2} \text{ sapwood s}^{-1}$ )









SPM2 site water balance components







Fig. 2. Site water balance components of the SPM2 model.

# Significance of winter C gain in southern pine systems










Martin 2000


# Air Temperature (C)











*Pinus elliottii*, Alachua County, FL, 1999-2001











- Drought strongly impacted C exchange
  - stomatal limitations dominant
  - longer term impacts of LAI and foliar [N]?
  - role of stand development?
- Freeze and cold temperatures strongly impacted C exchange
  - direct effects on  $V_{cmax}$
  - damage to PSII
  - stomatal response to VPD
- Nutrient limitations may be important
  - interaction of management, stand development, climate



# Further Questions

- To what extent does LAI influence drought responses?
  - Within-stand, temporal variation
  - Between-stand, ontogenetic variation
- Prevalence of soil drought
  - water table dependent?
- Interactions among temperature, water, nutrients?

# Acknowledgements

- Biological and Environmental Research Program (BER), U.S. Department of Energy, through the Southeast Regional Center of the National Institute for Global Environmental Change (NIGEC) under Cooperative Agreement No. DE-FC03-90ER61010.
- Forest Biology Research Cooperative
- Rayonier, Inc.
- David Nolletti
- Sean Gallagher